

A NOTE ON NON-SUPERSYMMETRIC OPEN-STRING ORBIFOLDS WITH A QUANTISED B_{AB} *

Carlo Angelantonj

Centre de Physique Théorique, École Polytechnique

F-91128 Palaiseau Cedex

and

Laboratoire de Physique Théorique de l'École Normale Supérieure

24 rue Lhomond, F-75231 Paris Cedex 05

angelant@cpt.polytechnique.fr

Abstract In this short note we review the main features of open-string orbifolds with a quantised flux for the NS-NS antisymmetric tensor in the context of the open descendants of non-supersymmetric asymmetric orbifolds with a vanishing cosmological constant.

Recently we have analysed in some detail the complete structure of the one-loop amplitudes for open strings [1, 2] on orbifolds [3] with a quantised NS-NS antisymmetric tensor background [4], thus extending the results in [5]. Besides recovering the expected rank reduction for the Chan-Paton (CP) gauge group, we were led to identify the discrete Wilson lines of [6] with the signs γ_ϵ that enforce a correct normalisation of the Möbius amplitude. In particular, for Z_2 orbifolds, they allow one to connect continuously $U(n)$ groups to $Sp(n) \otimes Sp(n)$ groups. In this brief note we want to reconsider the non-supersymmetric type I vacua with vanishing cosmological constant discussed in [7, 8] in the spirit of [4]. We refer to the original papers for more details and for references.

Non-supersymmetric vacua with a vanishing cosmological constant can be obtained as asymmetric orbifolds of type II and/or type I superstrings. The simplest instance of these models can be obtained consid-

*Preprint CPHT-PC231.0899, LPTENS 99/30

ering the generators [9]

$$\begin{aligned} f &= [(-1^4, 1; 1^5), (0^4, v_L; \delta^4, v_R), (-1)^{F_R}], \\ g &= [(1^5; -1^4, 1), (\delta^4, w_L; 0^4, w_R), (-1)^{F_L}]. \end{aligned} \quad (1.1)$$

The first entry in the square brackets denotes rotations, the second denotes shifts on the internal compactification lattice while the third corresponds to the space-time fermion-number. The shift δ acts as a Z_2 -shift, whereas $v_{L,R}$ ($= w_{R,L}$) act as an A_2 shift [10] on the fifth coordinate. The asymmetric nature of the orbifold together with level matching, require that the internal 4d lattice split into a product of four circles at the radius of SU(2) enhanced symmetry. Actually, the algebra of the generators (1.1) reveals that f and g generate a non-abelian space group orbifold S [11]. The restriction to the point group $\overline{P} = S/\{f^2, g^2\}$, with f^2 and g^2 pure translations, effectively changes the lattice to an SO(8) lattice, thus introducing a non-vanishing background for the B -field

$$B = \frac{\alpha'}{2} \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & -1 & -1 \\ 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}$$

with rank $r = 2$.

We have already noticed in [8] how the choice of the internal SO(8) lattice is crucial in order to get a consistent result. In fact, the orbifold that we are considering can be thought as an f (or g) projection of the supersymmetric T^4/Z_2 orbifold generated by fg , with the T^4 corresponding to the SO(8) lattice. The resulting torus amplitude

$$\mathcal{T}_0 \sim |V_4 O_4|^2 + |S_4 S_4|^2 - (O_4 V_4)(\bar{C}_4 \bar{C}_4) - (C_4 C_4)(\bar{O}_4 \bar{V}_4) + \dots$$

is then consistent with the generic Klein bottle amplitude¹ (that, roughly speaking feels only the left-right symmetric generator fg) [4]

$$\tilde{\mathcal{K}}_0 \sim \left[\sqrt{v} + \frac{2^{-r/2}}{\sqrt{v}} \right]^2 (V_4 O_4 - S_4 S_4) + \left[\sqrt{v} - \frac{2^{-r/2}}{\sqrt{v}} \right]^2 (O_4 V_4 - C_4 C_4)$$

only for an internal SO(8) lattice, where the volume $v = \frac{1}{2}$ and $r = 2$. Thus $\tilde{\mathcal{K}}_0$ contains only those states that in $\mathcal{T}_0$ are combined with their antiholomorphic counterparts. The same phenomenon also presents itself in the open sector, upon identification of Neumann and Dirichlet

¹In our case the spinors have a flipped chirality, since the supersymmetric Z_2 generator fg contains $(-1)^{F_L + F_R}$ together with the standard inversions.

CP charges as a result of modding out by the T-duality contained in f . As a result of the action of f , the CP gauge group is reduced further and turns into in a single U(8) factor [7, 8]. Actually, this is the case if we break the residual internal global $\text{SO}(4)^2$ symmetry by introducing discrete Wilson lines in the Möbius amplitude, thus affecting the $P = T^{\frac{1}{2}}ST^2ST^{\frac{1}{2}}$ transformation, that acts on the real hatted characters [6, 4]. Let us analyse the massless contribution to the transverse-channel Möbius amplitude:

$$\tilde{\mathcal{M}}_0 \sim -\frac{2}{2}(M_1 + M_2)[(\hat{V}_4\hat{O}_4 - \hat{S}_4\hat{S}_4)\hat{O}_4\hat{O}_4 - (\hat{O}_4\hat{V}_4 - \hat{C}_4\hat{C}_4)\hat{V}_4\hat{V}_4],$$

where with have explicitly written the contribution of the internal bosons. On the $\text{SO}(4)$ (hatted) characters $(\hat{O}_4, \hat{V}_4, \hat{S}_4, \hat{C}_4)$, the P matrix acts as $\text{diag}(\sigma_1, \sigma_1)$, with σ_1 the first Pauli matrix. As a result, the space-time vector contributes to

$$\mathcal{M}_0 \sim \frac{1}{2}(M_1 + M_2)[(\hat{V}_4\hat{O}_4 - \hat{S}_4\hat{S}_4)\hat{O}_4\hat{O}_4 - (\hat{O}_4\hat{V}_4 - \hat{C}_4\hat{C}_4)\hat{V}_4\hat{V}_4]$$

thus calling for a product of symplectic gauge groups $\text{Sp}(8) \otimes \text{Sp}(8)$, consistently with the tadpole conditions. The resulting model is still supersymmetric at all mass levels in the open sector, while it is non-supersymmetric in the closed sector. This has to be contrasted with the recently proposed scenario [12, 4] where supersymmetry is unbroken in the bulk (at any mass level) but is broken on branes, where a positive cosmological constant is generated at one loop.

The U(8) model of [7, 8] is obtained introducing in $\tilde{\mathcal{M}}$ discrete Wilson lines, that result in primed $\text{SO}(4)$ characters associated to the internal lattice [6, 4]:

$$\begin{aligned} \hat{O}_4 &= \hat{O}_2\hat{O}_2 - \hat{V}_2\hat{V}_2 \quad \rightarrow \quad \hat{O}'_4 = \hat{O}_2\hat{O}_2 + \hat{V}_2\hat{V}_2, \\ \hat{V}_4 &= \hat{O}_2\hat{V}_2 + \hat{V}_2\hat{O}_2 \quad \rightarrow \quad \hat{V}'_4 = \hat{O}_2\hat{V}_2 - \hat{V}_2\hat{O}_2, \end{aligned}$$

whose P transformation is now given by the Pauli matrix σ_3 acting on $(\hat{O}'_4, \hat{V}'_4)$. As a result

$$\mathcal{M}_0 \sim -\frac{1}{2}(M + \bar{M})[(\hat{O}_4\hat{V}_4 - \hat{C}_4\hat{C}_4)\hat{O}'_4\hat{O}'_4 - (\hat{V}_4\hat{O}_4 - \hat{S}_4\hat{S}_4)\hat{V}'_4\hat{V}'_4],$$

consistently with a unitary CP gauge group.

Although this latter choice has a sensible “geometric” 6d decompactification limit, it is not naturally related to the rational $\text{SO}(8)$ internal lattice. In fact, it is the $\text{Sp}(8) \otimes \text{Sp}(8)$ model that, in the 6d decompactification limit, leads to the $\text{Sp}(8)^4$ model of [2, 6, 13] pertaining to the rational $\text{SO}(8)$ lattice.

Acknowledgments

It is a pleasure to thank I. Antoniadis, K. Förger and A. Sagnotti for interesting discussions. I would also like to thank the Organizers of the Cargèse Summer School for having created a pleasant and stimulating atmosphere. This work was supported in part by EEC TMR contract ERBFMRX-CT96-0090.

References

- [1] A. Sagnotti, in Cargèse 87, Non-Perturbative Quantum Field Theory, eds. G. Mack et al. (Pergamon Press Oxford, 1988), p.521.
- [2] M. Bianchi and A. Sagnotti, Phys. Lett. B247 (1990) 517.
- [3] G. Pradisi and A. Sagnotti, Phys. Lett. B216 (1989) 59;
E. Gimon and J. Polchinski, Phys. Rev. D54 (1996) 1667.
- [4] C. Angelantonj, *Comments on open string orbifolds with a non-vanishing B_{ab}* , hep-th/9908064.
- [5] M. Bianchi, G. Pradisi and A. Sagnotti, Nucl. Phys. B376 (1992) 365.
M. Bianchi, Nucl. Phys. B528 (1998) 73;
E. Witten, JHEP 02 (1998) 006;
Z. Kakushadze, G. Shiu and S.-H.H. Tye, Phys. Rev. D58 (1998)
- [6] M. Bianchi and A. Sagnotti, Nucl. Phys. B361 (1991) 519.
- [7] R. Blumenhagen and L. Görlich, *Orientifolds of non-supersymmetric asymmetric orbifolds*, hep-th/9812158.
- [8] C. Angelantonj, I. Antoniadis and K. Förger, *Non-supersymmetric type I strings with zero vacuum energy*, hep-th/9904092;
K. Förger, these proceedings.
- [9] S. Kachru, J. Kumar and E. Silverstein, Phys. Rev. D59 (1999) 106004;
J.A. Harvey, Phys. Rev. D59 (1999) 26002.
- [10] C. Vafa and E. Witten, Nucl. Phys. Proc. Suppl. 46 (1996) 225.
- [11] L. Dixon, J.A. Harvey, C. Vafa and E. Witten, Nucl. Phys. B261 (1985) 678; Nucl. Phys. B274 (1986) 285.
- [12] I. Antoniadis, E. Dudas and A. Sagnotti, *Brane supersymmetry breaking*, hep-th/9908023;
G. Aldazabal and A.M. Uranga, *Tachyon-free non-supersymmetric type IIB orientifolds via brane-antibrane systems*, hep-th/9908072.
- [13] C. Angelantonj, M. Bianchi, G. Pradisi, A. Sagnotti and Ya.S. Stanev, Phys. Lett. B387 (1996) 743.